


Moesin; Clone MSN491 (Concentrate)


Availability/Contents:

<u>Item #</u>	<u>Volume</u>
RA0211-C.5	0.5 ml

Description:


Species: Mouse
 Immunogen: Recombinant human moesin protein
 Clone: MSN491
 Isotype: IgG1, kappa
 Entrez Gene ID: 4478 (Human)
 Hu Chromosome Loc.: Xq11.1
 Synonyms: Membrane-organizing extension spike protein; Moesin/anaplastic lymphoma kinase fusion protein; MSN/ALK fusion
 Mol. Weight of Antigen: 78kDa
 Format: 200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.
 Specificity: This antibody recognizes the 78kDa moesin protein.
 Background: Moesin, a member of the talin-4.1 superfamily, is a linking protein of the submembraneous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophage, lymphocyte, fibroblastic, endothelial, epithelial, and neuronal cell lines, but not in blood cells. The ERM proteins (ezrin, radixin, and moesin) are involved in a variety of cellular functions such as cell adhesion, migration, and the organization of cell surface structures. They are highly homologous both in protein sequence and in functional activity with merlin/schwannomin, a neurofibromatosis-2-associated tumor-suppressor protein. Cell lines of epithelial and mesothelial origin contain both moesin and radixin, whereas cells of endothelial and lymphoid origin express moesin.
 Species Reactivity: Human. Others not known.
 Positive Control: HT-29, CH3LC, or HUVEC cells. Uterus, placenta, tonsil (both B- and T-lymphocytes), skeletal muscle, thyroid, or kidney.
 Cellular Localization: Cell surface
 Titer/ Working Dilution: Immunohistochemistry (Frozen and Formalin-fixed): 0.5-1 µg/ml
 Flow Cytometry: 0.5-1 µg/million cells
 Immunofluorescence: 0.5-1 µg/ml
 Western Blotting: 0.5-1 µg/ml
 Immunoprecipitation: 0.5-1 µg/500µg protein lysate
 Microbiological State: This product is not sterile.

Storage: 2° C  8° C




ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.



 EmergoEurope (31)(0) 70 345-8570
Molsnstraat 15
2513 BH Hague, The Netherlands

Uses/Limitations: Not to be taken internally.
 For Research Use Only.
 This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.
 Do not use if reagent becomes cloudy.
 Do not use past expiration date.
 Non-Sterile.



Ordering Information and Current Pricing at www.scytek.com

Formalin-fixed, paraffin-embedded human placenta stained with Moesin; Clone MSN491.

Procedure:

1. **Tissue Section Pretreatment (Required):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).

Precautions: Contains Sodium Azide as a preservative (0.09% w/v).
 Do not pipette by mouth.
 Avoid contact of reagents and specimens with skin and mucous membranes.
 Avoid microbial contamination of reagents or increased nonspecific staining may occur.
 This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


References:

1. Lankes W *et. al.*, Biochem Journal, 1988; 251:831-842.
2. Schwartz-Albiez R *et. al.*, European Journal Cell Biology, 1995; 67:189-198.


Warranty:

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
 205 South 600 West
 Logan, UT 84321
 U.S.A.



 EmergoEurope (31)(0) 70 345-8570
 Molsnstraat 15
 2513 BH Hague, The Netherlands