

# TIMP-3 (Tissue Inhibitor of Metalloproteinase-3); Rabbit Polyclonal (Concentrate)

**Availability/Contents:**

<u>Item #</u>	<u>Volume</u>
RA0309-C.5	0.5 ml

**Description:**

**Species:** Rabbit

**Immunogen:** Recombinant fragment corresponding to Human TIMP3 aa 175-211

**Clone:** Rabbit polyclonal

**Isotype:** IgG

**Entrez Gene ID:** 7078 (Human); 21859 (Mouse); 25358 (Rat)

**Hu Chromosome Loc.:** 22q12.3

**Synonyms:** HSMRK222, K222TA2, Protein MIG-5, SFD, Sorsby fundus dystrophy pseudo-inflammatory, TIMP metalloproteinase inhibitor 3, Tissue Inhibitor of Metalloproteinase 3.

**Mol. Weight of Antigen:** 24kDa

**Format:** 200µg/ml of Ab purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.

**Specificity:** The amino acid sequence (a a175-211) used as the immunogen for anti-TIMP3 is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat.

**Background:** TIMP3 (tissue inhibitor of metalloproteinases 3), along with family members TIMP1, TIMP2, and TIMP4, are inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix (ECM). An imbalance between MMPs and the associated TIMPs may play a significant role in the invasive phenotype of malignant tumors. TIMPs inhibit the proteolytic invasiveness of tumor cells and normal placental trophoblast cells. TIMP-3 may be involved in regulating trophoblastic invasion of the uterus as well as in regulating remodeling of the extracellular matrix during the folding of epithelia, and in the formation, branching, and expansion of epithelial tubes.


**Species Reactivity:** Human, Mouse, Rat, Horse, Cow, and Dog. Others not known.

**Positive Control:** Placenta or breast carcinoma.

**Cellular Localization:** Cytoplasmic

**Titer/ Working Dilution:** Immunohistochemistry (Frozen and Formalin-fixed): 1-2 µg/ml  
 Flow Cytometry: 0.5-1 µg/million cells  
 Immunofluorescence: 1-2 µg/ml  
 Western Blotting: 0.5-1 µg/ml  
 Immunoprecipitation: 1-2 µg/500µg protein lysate

**Microbiological State:** This product is not sterile.

Storage: 2° C  8° C




ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

CE

 EmergoEurope (31)(0) 70 345-8570  
Molsnstraat 15  
2513 BH Hague, The Netherlands

**Uses/Limitations:** Not to be taken internally.  
For Research Use Only.  
This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.  
Do not use if reagent becomes cloudy.  
Do not use past expiration date.  
Non-Sterile.



Formalin-fixed, paraffin-embedded human placenta stained with TIMP-3; Rabbit Polyclonal.

**Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)**

**Procedure:**

1. **Tissue Section Pretreatment (Required):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).


**Precautions:** Contains Sodium Azide as a preservative (0.09% w/v).  
Do not pipette by mouth.  
Avoid contact of reagents and specimens with skin and mucous membranes.  
Avoid microbial contamination of reagents or increased nonspecific staining may occur.  
This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


**References:**

1. Nagase, H. et al. (2006) Cardiovasc Res 69, 562-73.
2. Visse, R. and Nagase, H. (2003) Circ Res 92, 827-39.

**Warranty:**

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.



 EmergoEurope (31)(0) 70 345-8570  
Molsnstraat 15  
2513 BH Hague, The Netherlands