

SOX10; Clone SOX10/991 (Concentrate)

Catalog Number

 A00161-C.1
 A00161-C

Volume

 0.1 ml
 1 ml

Description

Species: Mouse

Immunogen: Recombinant human SOX10 protein fragment (aa115-269) (exact sequence is proprietary).

Clone: SOX10/991

Isotype: IgG2b, Kappa.

Format: 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.

Specificity: Recognizes a protein of ~55kDa, identified as SOX10. This antibody does not cross-react with other members of the SOX-family.


Species Reactivity: Human and Mouse. Others-not known.

Positive Control: HepG2 cells, Melanoma, breast carcinomas, gliomas.

Cellular Localization: Nuclear

Titer/Working Dilution: Immunohistochemistry (Formalin-Fixed): 0.5-1 µg/ml

Microbiological State: Nonsterile.



Human melanoma stained using SOX10; Clone SOX10/991. Results were visualized using ScyTek's UltraTek Alk-Phos Anti-Mouse (Permanent Red) Staining System.

Intended Use

For In Vitro Diagnostic use. This antibody is intended for the qualitative visualization of the anatomical elements listed in the Specificity section. It is intended to be used within an Immunohistochemistry (IHC) procedure on formalin-fixed paraffin-embedded (FFPE) human tissue followed by visualization by light microscopy. Any diagnostic interpretation of the results of this antibody is to be complemented by morphological studies using proper controls and should be evaluated within the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Procedure

- Tissue Section Pretreatment (Required):** Staining of formalin fixed, paraffin embedded tissue sections is enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
- Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
- Visualization:** For maximum staining intensity we recommend ScyTek's UltraTek Alk-Phos Anti-Mouse (Permanent Red) Staining System (ScyTek Cat# AET080). See IFU for instructions.

Materials and Reagents Required but not Provided

- Control tissue and reagents.
- Xylene, graded alcohols, and deionized/distilled water.
- IHC detection system. Suggested: ScyTek's UltraTek Alk-Phos Anti-Mouse (Permanent Red) Staining System (ScyTek Cat# AET080).
- Wash buffer for rinses (ScyTek Cat# TBT500).
- Retrieval solution (ScyTek Cat# CPL500).
- Hematoxylin counterstain and bluing reagent (ScyTek Cat# HMM500 and BRT500).
- Mounting medium and coverslip.

Note: ScyTek Laboratories has a wide range of IHC reagents and ancillaries that can be found at scytex.com.

Storage and Stability

Do not Freeze. Store at 2-8°C. Return to 2-8° immediately after use. Do not use after expiration date printed on label. Verify visually that antibody has not been contaminated before use. Do not use if reagent becomes cloudy or precipitates.


Limitations

Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous alkaline phosphatase activity or endogenous biotin may cause non-specific staining depending on detection system used. This data sheet's recommendations and procedures were validated using ScyTek IHC reagents and may not be suitable for other detection systems.


Precautions

- Contains Sodium Azide as a preservative (0.09% w/v), do not ingest. Sodium Azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.
- Do not pipette by mouth.
- Avoid contact of reagents and specimens with skin and mucous membranes.
- Avoid microbial contamination of reagents or increased nonspecific staining may occur.

 Storage: 2° C  8° C



ScyTek Laboratories, Inc.
 205 South 600 West
 Logan, UT 84321
 U.S.A.

Emergo Europe
 Prinsessegracht 20
 2514 AP The Hague, The Netherlands

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 – Tel. (435) 755-9848 - Fax (435) 755-0015 - www.ScyTek.com


5. The user must validate any procedures and recommendations that differ from this data sheet.
6. The SDS may be found at scytex.com


References

1. Mohamed A, et al. SOX10 Expression in malignant melanoma, carcinoma, and normal tissues. Appl Immunohistochem Mol Morphol. 2013; 21(6):506-10.

Warranty

No products or "Instructions For Use (IFU)" are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE 

EC REP

Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands