

Instructions For Use ACU-IFU

Rev. Date 4/22/2019

Revision: 1

Page 1 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Fax (435) 755-0015 - www.scytek.com


DAB Substrate (High Contrast)

Description:

DAB Substrate (High Contrast) is a solution specifically designed for optimal signal intensity and contrast. When in the presence of peroxidase enzyme and DAB Chromogen, the substrate helps create a brown precipitate that is insoluble in alcohol. The standard working dilution is 50ul (0.9mg) of DAB Chromogen per 1ml of DAB Substrate (High Contrast), although the ratio can be adjusted as desired, the use of liquid components reduces some risks associated with handling powders (ie. dust inhalation), and eliminates waste which often results from using tablets that require a predetermined final volume. Once the two components are combined, the reagent can be used for up to six hours, making it ideal for automated stainers.

Uses/Limitations:

Not to be taken internally.
For In-Vitro Diagnostic Use only.
Hematologic applications.
Do not use if reagents become cloudy.
Do not use past expiration date.
Use caution when handling reagents.
Non-Sterile.



Lung Squamous Cell Carcinoma Stained with ScyTek's Cytokeratin 5; Clone EP42 (Cat#A00139) and an IHC protocol containing ScyTek's DAB Substrate (ACU)

Control Tissue:

Any well-fixed frozen or FFPE tissue.

Availability:

Item #	Volume
ACU250	250ml
ACU500	500ml
ACU999	1000ml

Storage:


Store at 2-8°C. Stable for 18 months from the date of manufacture.


Ordering Information and Current Pricing at www.scytek.com

Precautions:

DAB Chromogen: Contains Diaminobenzidine in buffer. DAB is a suspected carcinogen.. Avoid contact with skin and eyes. Reagent is acidic and can cause burns if skin contact occurs. Handle with care and dispose of according to regulations.

DAB Substrate: Contains Hydrogen Peroxide in buffer. Avoid contact with skin and eyes.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE IVD

EC REP
Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands

Instructions For Use
ACU-IFU

Rev. Date 4/22/2019

Revision: 1

Page 2 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Fax (435) 755-0015 - www.scytek.com

Required but not Provided:


DAB Chromogen (ScyTek Cat#: ACB)


Procedure:

1. After incubation and rinse of HRP, Combine 50µl of DAB Chromogen with 1mL of DAB Substrate (High Contrast).
2. Apply mixture to tissue section.
3. Incubate tissue section for 5-15 minutes or until desired reaction intensity.
4. Rinse tissue in buffer, counterstain, dehydrate, and clear in xylene or xylene substitute and coverslip.

References:

1. Keiichi Abe, Ryo Shimada, Yoshikazu Okada & Kazuhiko Kibayashi (2016) Traumatic brain injury decreases serotonin transporter expression in the rat cerebrum, *Neurological Research*, 38:4, 358-363, DOI: 10.1080/01616412.2015.1110402
2. Tominaga T, Shimada R, Okada Y, Kawamata T, Kibayashi K (2019) Senescence-associated-β-galactosidase staining following traumatic brain injury in the mouse cerebrum. *PLoS ONE* 14(3): e0213673. <https://doi.org/10.1371/journal.pone.0213673>
3. Furat Rençber, Selenay, Sema Kurnaz Ozbek, Ceyla Eraldemir, Zehra Sezer, Tugba Kum, Sureyya Ceylan, and Elif Guzel. "Effect of Resveratrol and Metformin on Ovarian Reserve and Ultrastructure in PCOS: An Experimental Study." *Journal of Ovarian Research* 11, no. 1 (June 29, 2018): 55. <https://doi.org/10.1186/s13048-018-0427-7>.
4. Georgiev, Georgi P, Boycho Landzhov, Iva N Dimitrova, Svetoslav Slavchev, Lina Malinova, Yuliyann Kartelov, Dessislava Ankova, and Wladimir Ovtsharov. "LIGHT MICROSCOPIC AND IMMUNOHISTOCHEMICAL STUDY OF THE MEDIAL COLLATERAL LIGAMENT EPILIGAMENT IN RAT KNEE," n.d., 8.
5. Georgiev, Georgi P, Boycho Landzhov, Iva N Dimitrova, Svetoslav Slavchev, Lina Malinova, and Wladimir Ovtsharov. "IMMUNOHISTOCHEMICAL STUDY DURING EARLY HEALING OF THE MEDIAL COLLATERAL LIGAMENT EPILIGAMENT IN RAT KNEE MODEL," n.d., 8.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE 

EC REP
Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands