

Instructions For Use AEX080-IFU

Revision: 3

Rev. Date: Nov. 4, 2021

Page 1 of 3

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

EconoTek HRP Anti-Polyvalent (DAB) Ready-To-Use (70 slide)

Description:	Species of Origin: Antigen Specificity: Preadsorbed Against: Enzyme Conjugate: Chromogen Substrate:	Goat Anti-Mouse, Rat, Rabbit, Guinea Pig Human Peroxidase Diaminobenzidine (DAB)
Contents:	Super Block Peroxide Block EconoTek Anti-Polyvalent EconoTek HRP DAB Chromogen DAB Substrate	8 ml 8 ml 8 ml 8 ml 3 ml 5 ml x 7 vials
Uses/Limitations:	Not to be taken internally. For In-Vitro Diagnostic use only. Do not use if reagent becomes cloudy. Do not use past expiration date. Use caution when handling reagents. Non-Sterile.	
Storage:	Store at 2-8°C.	
Precautions:	Avoid contact with skin and eyes. Harmful if swallowed. Follow all Federal, State, and local regulations regarding disposal.	

Procedure:

- 1. Deparaffinize and rehydrate tissue section.
- 2. To reduce nonspecific background staining due to endogenous peroxidase, incubate slide in hydrogen peroxide for 5 minutes.
- 3. Wash 2 times in buffer.
- 4. If required, incubate tissue in digestive enzyme.
- 5. Wash 4 times in buffer.
- 6. Apply Super Block (blue cap), and incubate for 5-10 minutes at room temperature to block nonspecific background staining. Note: Do not exceed 10 minutes or there may be a reduction in desired stain.
- 7. Wash 1 time in buffer.
- 8. Apply primary antibody and incubate according to manufacturer's protocol.

Storage: 2° C

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.



Emergo Europe Prinsessegracht 20 2514 AP The Hague, The Netherlands



Instructions For Use AEX080-IFU

Revision: 3

Rev. Date: Nov. 4, 2021

Page 2 of 3

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

- 9. Wash 4 times in buffer.
- 10. Apply EconoTek Biotinylated Anti-polyvalent (yellow cap), and incubate for 30 minutes at room temperature.
- 11. Wash 4 times in buffer.
- 12. Apply EconoTek HRP (red cap), and incubate for 30 minutes at room temperature.
- 13. Rinse 4 times in buffer.
- 14. Add 8 drops of DAB chromogen to one 5ml vial of DAB Substrate. Mix well and apply to tissue for 5 minutes.
- 15. Rinse 1 time in Deionized water.
- 16. Apply chromogen/substrate mixture to tissue for another 5 minutes.
- 17. Rinse in Deionized water.
- 18. Counterstain as desired.

WARNING:DAB is a suspected carcinogen. Handle with care and dispose of according to all regulations.

19. Counterstain and coverslip

Troubleshooting Guide

Overstaining:

- 1. Concentration of the primary antibody was too high or the incubation time was too long.
- 2. Temperature during incubation was too high.
- 3. Incubation time with link antibody or streptavidin/enzyme label was too long.

Nonspecific Background Staining:

- 1. Inadequate rinsing between steps.
- 2. Tissue was allowed to dry with reagents on.
- 3. Folds in tissue trapped reagents.
- 4. Tissue contains endogenous peroxidase.
- 5. Tissue contains endogenous biotin.
- 6. Antigen migrated in tissue.
- 7. Excessive tissue adhesive on slides.
- 8. Inadequate blocking with protein block.

Weak Staining:

- 1. Primary antibody concentration was too low or incubation time was too short.
- 2. Reagents are past their expiration date.
- 3. Inadequate removal of wash water between steps, resulting in dilution of reagents.
- 4. Room temperature was excessively cool.
- 5. The primary antibody does not recognize an antigen that survives fixation and embedding in high enough amounts.
- 6. Excessive incubation with protein block (Super Block).

No Staining:

1. Steps were inadvertently left out.

8° C Storage: 2° C

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.



Emergo Europe Prinsessegracht 20 2514 AP The Hague, The Netherlands



Instructions For Use EX080-IFU А **Revision: 3**

Rev. Date: Nov. 4, 2021

Page 3 of 3

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

- 2. There is no antigen in the tissue.
- The primary antibody is not of mouse, rat, rabbit or guinea pig origin. 3.
- 4. Chromogenic substrate has been replaced with another that is not intended for use with peroxidase.







EC REP Emergo Europe Prinsessegracht 20 2514 AP The Hague, The Netherlands