

Anti-mouse CD31 / DIA-310

Rat monoclonal anti-mouse endothelial cell marker CD31 (PECAM-1), Clone SZ31

Product Information

Catalog No.:	DIA-310 (100µg)	Reconstitution:	DIA-310 (100 µg), restore to 500 µl
Clone:	SZ31		Reconstitute with sterile distilled water
Concentration:	0.2 mg/ml	Presentation:	by gentle shaking for 10 minutes
Isotype:	Rat IgG2a		PBS with 2% BSA, 0.05% NaN ₃ , pH 7.4.
Specificity:	Murine CD31 (PECAM-1) (adult and embryonic endothelial cells)	Applications:	Antibody purified from culture supernatant
Immunogen:	Murine amino acid fragment		Immunohistochemistry (standard formalin-fixed paraffin and frozen sections)
Species:	Mouse, pig,	Dilutions:	Western blot
Reactivity:	does not cross-react with rat or human		1:20 Immunohistochemistry (IHC)
Physical State:	Lyophilized powder		1:5000 Western Blot
			(General recommendation, validation of antibody performance/protocol using proper controls is the responsibility of the end user.)

Reactivity

Antibody clone SZ31 is the first antibody which reacts specifically with murine CD31 in formalin-fixed paraffin-embedded tissue sections.

CD31, also known as PECAM-1 (Platelet Endothelial Cell Adhesion Molecule-1) is expressed constitutively on the surface of embryonic and adult endothelial cells. It is also expressed on cell surfaces of monocytes, neutrophils, platelets and certain T-cell subsets. It has been detected on bone marrow-derived hematopoietic stem cells and embryonic stem cells. CD31 is a 130kDa integral membrane glycoprotein and as a member of the immunoglobulin superfamily involved in the mediation of cell-to-cell adhesion. CD31-mediated endothelial cell-cell interactions play a major role in angiogenesis. Studies have shown CD31 to be a superior marker in human angiogenesis, which reportedly predicts tumor recurrence. Pathophysiological studies of CD31 in murine model systems had limitations because standard formalin-fixed sections were excluded. The clone SZ31 eliminates these restrictions by allowing high quality immunohistochemical (IHC) analysis of standard formalin-fixed paraffin sections in mice.

Instructions for Use

Immunohistochemical staining of standard formalin-fixed paraffin sections

Indirect alkaline phosphatase staining (Other techniques, e.g. Avidin-Biotin-alkaline phosphatase (ABAP), alkaline phosphatase anti-alkaline phosphatase (AAPAP) or horseradish peroxidase (HRP) -method are also possible).

1. Deparaffinize formalin-fixed paraffin-embedded mouse tissue sections by a standard procedure using xylol/ethanol
2. Antigen retrieval: high temperature heating of sections in citrate buffer pH 6,0 according to standard procedures
3. Block with 5% rabbit serum, 10 min RT
4. Wash with TBS, 3 x 5 min
5. Incubate with DIA-310 (1:10-1:20), 30min RT
6. Wash with TBS, 3 x 5 min
7. Incubate with rabbit anti-rat IgG (H+L) alkaline phosphatase (1:200), 30min RT
8. Wash with TBS, 3 x 5 min
9. Add substrate, e.g. Neufuchsin, 30min RT
10. Counterstain, e.g. with Hematoxylin-Papanicolaou

Storage and Stability

The antibody clone SZ31 in lyophilised form is stable for at least one year (-20°). As reconstituted liquid store at 2-8°C short term (several weeks). For long term storage aliquot and freeze at -20°C or -80°C. Avoid repeated freeze / thaw cycles

Safety Notes

The material contains 0.05% sodium azide as preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material. Avoid skin and eye contact, inhalation, and ingestion.




Figure 1

Immunohistochemistry of mouse CD31 (PECAM-1) in formalin-fixed paraffin-embedded tissue sections

(pictures courtesy of Prof. Dr. Robert Klopffleisch, Institute of Pathology, Department of Veterinary Pathology, Berlin, Germany)

The monoclonal antibody clone SZ31 reacts specifically with endothelial cells in vessels and capillaries of aorta (A), aortic origin (B), endocardium (C), small intestine (D, E), Colon (F), brain (G), lymph nodes (H), bone marrow (I), mesenteric vessels (J), kidney (K) and pancreas (L). All sections were stained by an indirect horseradish peroxidase (HRP)-method according to standard procedures, counterstaining with Hämatoxylin.



For research use only. Not for diagnostic or therapeutic use.



Dianova GmbH
Warburgstrasse 45
D-20354 Hamburg
Germany

URL: www.dianova.com
Email: info@dianova.com
Phone: +49 (0)40 – 45067 – 0
Fax: +49 (0)40 – 45067 – 490

dianova

Figure 2

Immunohistochemistry of mouse CD31 (PECAM-1) in formalin-fixed paraffin-embedded tissue sections

(pictures courtesy of Prof. Dr. H. Stein, Institute of Pathology, Charité Campus Benjamin Franklin, Berlin, Germany)

The monoclonal antibody clone SZ31 reacts specifically with endothelial cells in vessels and capillaries of murine lung (A), skeletal muscle (B), spinal cord (C), liver (D), and murine adenocarcinoma (E, F). All sections were stained by an indirect alkaline phosphatase method according to standard procedures with antigen retrieval by high-temperature heating in citrate buffer and counterstaining with Hämatoxylin-Papanicolaou.





Figure 3

Western blot analysis: Immunoblot of extracts from murine lung, J558L cells and m-Lend cells using CD31 rat monoclonal antibody clone SZ31 (DIA-310 1:5.000) and goat anti-rat-HRP antibody (1:10.000)

Lane:	1	J558 cells (CD31-), 25µg lysate
	2	murine lung, 25µg lysate
	3	blank
	4	m-Lend cells (CD31+), 12.5µg lysate



For research use only. Not for diagnostic or therapeutic use.



Dianova GmbH
Warburgstrasse 45
D-20354 Hamburg
Germany

URL: www.dianova.com
Email: info@dianova.com
Phone: +49 (0)40 – 45067 – 0
Fax: +49 (0)40 – 45067 – 490



Selection of Specific References for anti-ms CD31, clone SZ31

1. Kim K, Watson PA, Lebda S, Jebiwott S, Somma A, La Rosa SP, Mehta D, Murray KS, Lilja HG, Ulmert D, Monette S, Scherz AJ, Coleman J. Androgen Deprivation Therapy Potentiates the Efficacy of Vascular Targeted Photodynamic Therapy of Prostate Cancer Xenografts. *Clin Cancer Res*. 2018 Feb 20; PMID: 29463549
2. de Jong RJ, Paulin N, Lemnitzer P, Viola JR, Winter C, Ferraro B, Grommes J, Weber C, Reutelingsperger C, Drechsler M, Soehnlein O. Protective Aptitude of Annexin A1 in Arterial Neointima Formation in Atherosclerosis-Prone Mice-Brief Report. *Arterioscler Thromb Vasc Biol*. 2017 Feb;37(2):312–315. PMID: 28062503
3. Looney AP, Han R, Stawski L, Marden G, Iwamoto M, Trojanowska M. Synergistic Role of Endothelial ERG and FLI1 in Mediating Pulmonary Vascular Homeostasis. *Am J Respir Cell Mol Biol*. 2017;57(1):121–131. PMID: PMC5516275
4. Matsuki M, Adachi Y, Ozawa Y, Kimura T, Hoshi T, Okamoto K, Tohyama O, Mitsuhashi K, Yamaguchi A, Matsui J, Funahashi Y. Targeting of tumor growth and angiogenesis underlies the enhanced antitumor activity of lenvatinib in combination with everolimus. *Cancer Sci*. 2017 Apr;108(4):763–771. PMID: PMC5406533
5. Yanagida K, Liu CH, Faraco G, Galvani S, Smith HK, Burg N, Anrather J, Sanchez T, Iadecola C, Hla T. Size-selective opening of the blood–brain barrier by targeting endothelial sphingosine 1–phosphate receptor 1. *PNAS*. 2017 Apr 25;114(17):4531–4536. PMID: 28396408
6. Zaccagnino A, Managò A, Leanza L, Gontarewitz A, Linder B, Azzolini M, Biasutto L, Zoratti M, Peruzzo R, Legler K, Trauzold A, Kalthoff H, Szabo I. Tumor-reducing effect of the clinically used drug clofazimine in a SCID mouse model of pancreatic ductal adenocarcinoma. *Oncotarget*. 2017 Jun 13;8(24):38276–38293. PMID: PMC5503532
7. Stout-Delgado HW, Cho SJ, Chu SG, Mitzel DN, Villalba J, El-Chemaly S, Ryter SW, Choi AMK, Rosas IO. Age-Dependent Susceptibility to Pulmonary Fibrosis Is Associated with NLRP3 Inflammasome Activation. *Am J Respir Cell Mol Biol*. 2016;55(2):252–263. PMID: PMC4979364
8. Goumas FA, Holmer R, Egberts J-H, Gontarewicz A, Heneweer C, Geisen U, Hauser C, Mende M-M, Legler K, Röcken C, Becker T, Waetzig GH, Rose-John S, Kalthoff H. Inhibition of IL-6 signaling significantly reduces primary tumor growth and recurrences in orthotopic xenograft models of pancreatic cancer. *Int J Cancer*. 2015 Sep 1;137(5):1035–1046. PMID: 25604508
9. Yue GG-L, Lee JK-M, Kwok H-F, Cheng L, Wong EC-W, Jiang L, Yu H, Leung H-W, Wong Y-L, Leung P-C, Fung K-P, Lau CB-S. Novel PI3K/AKT targeting anti-angiogenic activities of 4-vinylphenol, a new therapeutic potential of a well-known styrene metabolite. *Sci Rep*. 2015 Jun 8;5:11149. PMID: PMC4459151
10. Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat MM, Zeitz M, Siegmund B, Kühl AA. A guide to histomorphological evaluation of intestinal inflammation in mouse models. *Int J Clin Exp Pathol*. 2014;7(8):4557–4576. PMID: PMC4152019
11. Copeland BT, Bowman MJ, Ashman LK. Genetic ablation of the tetraspanin CD151 reduces spontaneous metastatic spread of prostate cancer in the TRAMP model. *Mol Cancer Res*. 2013 Jan;11(1):95–105. PMID: 23131993
12. Martin-Padura I, Marighetti P, Agliano A, Colombo F, Larzabal L, Redrado M, Bleau A-M, Prior C, Bertolini F, Calvo A. Residual dormant cancer stem-cell foci are responsible for tumor relapse after antiangiogenic metronomic therapy in hepatocellular carcinoma xenografts. *Lab Invest*. 2012 Jul;92(7):952–966. PMID: 22546866
13. Hölscher M, Silter M, Krull S, von Ahlen M, Hesse A, Schwartz P, Wielockx B, Breier G, Katschinski DM, Zieseniss A. Cardiomyocyte-specific prolyl-4-hydroxylase domain 2 knock out protects from acute myocardial ischemic injury. *J Biol Chem*. 2011 Apr 1;286(13):11185–11194. PMID: PMC3064173
14. Kröger C, Vijayaraj P, Reuter U, Windoffer R, Simmons D, Heukamp L, Leube R, Magin TM. Placental vasculogenesis is regulated by keratin-mediated hyperoxia in murine decidual tissues. *Am J Pathol*. 2011 Apr;178(4):1578–1590. PMID: PMC3078447
15. Agliano A, Martin-Padura I, Marighetti P, Gregato G, Calleri A, Prior C, Redrado M, Calvo A, Bertolini F. Therapeutic effect of lenalidomide in a novel xenograft mouse model of human blastic NK cell lymphoma/blastic plasmacytoid dendritic cell neoplasm. *Clin Cancer Res*. 2011 Oct 1;17(19):6163–6173. PMID: 21856771
16. Anania MC, Sensi M, Radaelli E, Miranda C, Vizioli MG, Pagliardini S, Favini E, Cleris L, Supino R, Formelli F, Borrello MG, Pierotti MA, Greco A. TIMP3 regulates migration, invasion and in vivo tumorigenicity of thyroid tumor cells. *Oncogene*. 2011 Jul 7;30(27):3011–3023. PMID: 21339735
17. Gramann M, Wendler O, Haeberle L, Schick B. Prominent collagen type VI expression in juvenile angiofibromas. *Histochem Cell Biol*. 2009 Jan;131(1):155–164. PMID: 18797915

Find more specific references for anti-ms CD31, clone SZ31 at www.dianova.com

For research use only. Not for diagnostic or therapeutic use.



Dianova GmbH
Warburgstrasse 45
D-20354 Hamburg
Germany

URL: www.dianova.com
Email: info@dianova.com
Phone: +49 (0)40 – 45067 – 0
Fax: +49 (0)40 – 45067 – 490

