

# HSP27 (Heat Shock Protein 27); Clone 24K/774 (Concentrate)

**Availability/Contents:**

<u>Item #</u>	<u>Volume</u>
RA0132-C.5	0.5 ml

**Description:**

**Species:** Mouse

**Immunogen:** Recombinant human HSPB1 protein.

**Clone:** 24K/774

**Isotype:** IgG1, kappa

**Entrez Gene ID:** 3315 (Human); 15507 (Mouse); 24471 (Rat)

**Hu Chromosome Loc.:** 7q11.23

**Synonyms:** Heat shock 27kDa protein, Estrogen-regulated 24kDa protein, Heat shock 25kDa protein 1, Heat shock 28kDa protein 1, Heat shock protein beta-1, HMN2B, HSP25, HSP28, HSPB1, Stress-responsive protein 27 (SRP27)

**Mol. Weight of Antigen:** 27kDa

**Format:** 200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.

**Specificity:** This antibody recognizes a 24-27kDa estrogen-regulated protein, identified as heat shock protein 27 (HSP27).

**Background:** HSP27 was recently found to be identical to the estrogen-induced “p29” and “24K” protein. About 50% of breast carcinomas are positive for HSP27, especially those that are also positive for estrogen and/or progesterone receptor. HSP27 has also been implicated in drug resistance in cancer cells.

**Species Reactivity:** Human, Chimpanzee, Monkey, Sheep, Rat, Mouse, and Chicken. Others not known.


**Positive Control:** ~50% of breast carcinomas are positive for HSP27, especially those that are also positive for estrogen and/or progesterone receptor.

**Cellular Localization:** Predominantly cytoplasmic with some nuclear.

**Titer/ Working Dilution:** Immunohistochemistry (Frozen and Formalin-fixed): 0.5-1 µg/ml  
 Flow Cytometry: 0.5-1 µg/million cells  
 Immunofluorescence: 0.5-1 µg/ml  
 Western Blotting: 0.25-0.5 µg/ml  
 Immunoprecipitation: 0.5-1 µg/500µg protein lysate

**Microbiological State:** This product is not sterile.

Storage: 2° C  8° C




ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

**CE**

 EmergoEurope (31)(0) 70 345-8570  
Molsnstraat 15  
2513 BH Hague, The Netherlands

**Uses/Limitations:** Not to be taken internally.  
 For Research Use Only.  
 This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.  
 Do not use if reagent becomes cloudy.  
 Do not use past expiration date.  
 Non-Sterile.



Formalin-paraffin breast carcinoma stained with HSP27; Clone 24K/774.

**Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)**

**Procedure:**

1. **Tissue Section Pretreatment (Highly Recommended):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).

**Precautions:** Contains Sodium Azide as a preservative (0.09% w/v).  
 Do not pipette by mouth.  
 Avoid contact of reagents and specimens with skin and mucous membranes.  
 Avoid microbial contamination of reagents or increased nonspecific staining may occur.  
 This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


**References:**

1. Georgopoulos, C. and Welch, W.J. 1993. Role of the major heat shock proteins as molecular chaperones. Annu. Rev. Cell Biol. 9: 601-634.

**Warranty:**

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.  
 205 South 600 West  
 Logan, UT 84321  
 U.S.A.



 EmergoEurope (31)(0) 70 345-8570  
 Molsnstraat 15  
 2513 BH Hague, The Netherlands