

Fascin-1 (Reed-Sternberg Cell Marker); Clone FSCN1/417 (Concentrate)

| | | |
|-------------------------------|---------------|---------------|
| Availability/Contents: | <u>Item #</u> | <u>Volume</u> |
| | RA0450-C.1 | 0.1 ml |
| | RA0450-C.5 | 0.5 ml |
| | RA0450-C1 | 1 ml |

Description:

| | |
|--------------------------|---|
| Species: | Mouse |
| Immunogen: | Full length recombinant human FSCN1 protein |
| Clone: | FSCN1/417 |
| Isotype: | IgG2a, kappa |
| Entrez Gene ID: | 6624 (Human) |
| Hu Chromosome Loc.: | 7p22.1 |
| Synonyms: | 55kDa actin-bundling protein; FAN1; Fascin homolog 1; Fascin1; FSCN1; HSN; p55; Singed (Drosophila) like (sea urchin fascin homolog like); SNL. |
| Mol. Weight of Antigen: | 55kDa |
| Format: | 200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. |
| Specificity: | This antibody recognizes a protein of 55kDa, which is identified as fascin-1. |
| Background: | The ability of fascin-1 to bind actin is regulated by phosphorylation. An antibody to fascin-1 is a very sensitive marker for Reed-Sternberg cells and variants in nodular sclerosis, mixed cellularity, and lymphocyte depletion Hodgkin's disease. It is uniformly negative in lymphoid cells, plasma cells, and myeloid cells. Fascin-1 is also expressed in dendritic cells. This marker may be helpful to distinguish between Hodgkin's lymphoma and non-Hodgkin's lymphoma in difficult cases. Also, the lack of expression of fascin-1 in the neoplastic follicles in follicular lymphoma may be helpful in distinguishing these lymphomas from reactive follicular hyperplasia in which the number of follicular dendritic cells is normal or increased. An antibody to fascin-1 has been suggested as a prognostic marker in neuroendocrine neoplasms of the lung as well as in ovarian cancer. Fascin-1 expression may be induced by Epstein-Barr virus (EBV) infection of B-cells with the possibility that viral induction of fascin in lymphoid or other cell types must also be considered in EBV-positive cases. |
| Species Reactivity: | Human. Others not known. |
| Positive Control: | HeLa, MCF-7, PC-3 or BEWO cells. Hodgkin's lymphoma, Ovarian, or Testicular Carcinoma. |
| Cellular Localization: | Cytoplasmic |
| Titer/ Working Dilution: | Immunohistochemistry (Frozen and Formalin-fixed): 0.5-1 µg/ml Flow Cytometry: 0.5-1 µg/million cells Immunofluorescence: 1-2 µg/ml Western Blotting: 0.5-1 µg/ml Immunoprecipitation: 1-2 µg/500µg protein lysate |
| Microbiological State: | This product is not sterile. |

Storage: 2° C  8° C




ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE

EC | REP
Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands

Uses/Limitations: Not to be taken internally.
 For Research Use Only.
 This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.
 Do not use if reagent becomes cloudy.
 Do not use past expiration date.
 Non-Sterile.



Ordering Information and Current Pricing at www.scytek.com

Formalin-fixed, paraffin-embedded human testicular carcinoma stained with Fascin-1; Clone FSCN1/417.

Procedure:

1. **Tissue Section Pretreatment (Required):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).

Precautions:


Contains Sodium Azide as a preservative (0.09% w/v).
 Do not pipette by mouth.
 Avoid contact of reagents and specimens with skin and mucous membranes.
 Avoid microbial contamination of reagents or increased nonspecific staining may occur.
 This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


References:

1. Yamashiro-Matsumura S and Matsumura F. J Biol Chem 1985; 260(8): 5087.
2. Yamashiro-Matsumura S and Matsumura F. J Cell Biol 1986; 103:631.
3. Duh F-M, et al. DNA Cell Biol 1994; 13(8):821.

Warranty:

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
 205 South 600 West
 Logan, UT 84321
 U.S.A.

CE

EC REP
 Emergo Europe
 Prinsessegracht 20
 2514 AP The Hague, The Netherlands