

CD100 (Semaphorin-4D); Clone 133-1C6 (Concentrate)

Availability/Contents:	<u>Item #</u>	<u>Volume</u>
	RA0492-C.1	0.1 ml
	RA0492-C.5	0.5 ml
	RA0492-C1	1 ml

Description:

Species:	Mouse
Immunogen:	PHA stimulated human peripheral blood lymphocytes.
Clone:	133-1C6
Isotype:	IgM, kappa
Entrez Gene ID:	10507
Hu Chromosome Loc.:	9q22.2
Synonyms:	COLL4; Collapsin4; GR3; M sema G; MSEMA; SEMA4D; Semacl2; SemaH; SEMAJ; Semaphorin C like 2; Semaphorin H; Semaphorin J; Semaphorin-4D; Semcl2
Mol. Weight of Antigen:	50kDa (Monomer)
Format:	200ug/ml of Ab purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA 0.05% azide.
Specificity:	Recognizes a homodimeric protein comprised of 50kDa subunits, identified as CD100 (Workshop VI; Code N-L026).
Background:	CD100 is expressed on majority of haemopoietic cells (B, T, NK and myeloid cells) and is absent from bone marrow, erythrocytes, eosinophils and endothelial cells. Its expression is increased after PHA-activation. CD100 was shown to associate with different partner molecules in T cells such as CD45, a key molecule with protein tyrosine phosphatase activity involved in T-cell transduction, and a Serine kinase. It plays a role in homotypic cell adhesion and in T cell activation.
Species Reactivity:	Human and Baboon. Does not react with horse. Others not-tested.
Positive Control:	Daudi, Raji, HUT-78, Kg1a, U937, and human lymphocytes. Human tonsil and lymph node.
Cellular Localization:	Cell surface.
Titer/ Working Dilution:	Flow Cytometry: 1-2 µg/million cells Immunofluorescence: 1-2 µg/ml Optimal dilution for a specific application should be determined by user.
Microbiological State:	This product is not sterile.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE
EC REP
Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands

Uses/Limitations: Not to be taken internally.
For Research Use Only.
Do not use if reagent becomes cloudy.
Do not use past expiration date.
Non-Sterile.

Ordering Information and Current Pricing at www.scytek.com

Procedure:



1. Optimal dilution and procedure for a specific application should be determined by user.

Precautions: Contains Sodium Azide as a preservative (0.09% w/v).
Do not pipette by mouth.
Avoid contact of reagents and specimens with skin and mucous membranes.
Avoid microbial contamination of reagents or increased nonspecific staining may occur.
This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.

References:

1. Kishimoto T et al. (eds) Leukocyte Typing VI, Garland Publishing, New York, 1997.
2. Schlossman SF et al. (eds) Leucocyte Typing V Oxford University Press, Oxford, 1995.
3. Delaire S et al. CD100 is a leukocyte semaphorin. Cell Mol Life Sci 1998, 54(11):1265-1276.
4. Dorfman DM et al. The leukocyte semaphorin CD100 is expressed in most T-cell, but few B-cell, non-Hodgkin's lymphomas. Am J Pathol 1998, 153(1):255-262.
5. Herold C et al. CD100 is associated with CD45 at the surface of human T lymphocytes. Role in T cell homotypic adhesion. J Immunol 1996, 157(12):5262-5268.
6. Elhabazi A et al. The human semaphorin-like leukocyte cell surface molecule CD100 associates with a serine kinase activity. J Biol Chem 1997, 272(38):23515-23520.
7. Hall KT et al. Human CD100, a novel leukocyte semaphorin that promotes B-cell aggregation and differentiation. Proc Natl Acad Sci USA 1996, 93(21):11780-11785

Warranty: No products or "Instructions For Use (IFU)" are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.**CE**

EC REP

Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands