


# CD26 (DPP IV / ADA-Binding Protein); Clone 134-2C2 (Concentrate)

<b>Availability/Contents:</b>	<u>Item #</u>	<u>Volume</u>
	RA0561-C.1	0.1 ml
	RA0561-C.5	0.5 ml
	RA0561-C1	1 ml

**Description:**

Species:	Mouse.
Immunogen:	A synthetic peptide from the amino terminal region of CD26.
Clone:	134-2C2.
Isotype:	IgM, kappa.
Entrez Gene ID:	1803.
Hu Chromosome Loc.:	2q24.3.
Synonyms:	ADA-binding protein (ADABP); Adenosine deaminase complexing protein 2 (ADCP-2); Dipeptidyl peptidase 4; Dipeptidyl peptidase 4 soluble form; Dipeptidyl peptidase IV membrane form; Dipeptidyl peptidase IV soluble form; Dipeptidyl peptidase, intestinal; Dipeptidylpeptidase 4; Dipeptidylpeptidase IV; DPP4; DPPIV; Intestinal dipeptidyl peptidase; T-cell activation antigen CD26; TP103.
Mol. Weight of Antigen:	110kDa.
Format:	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.
Specificity:	Recognizes a glycoprotein of 110kDa, identified as CD26.
Background:	It is an atypical serine protease belonging to the prolyl oligopeptidase family. It is expressed on lymphocyte cells and is upregulated during T-cell activation. CD26 is also expressed on activated B cells and natural killer cells and abundantly on epithelia. CD26 is implicated in a variety of biological functions including T-cell activation, cell adhesion with extracellular matrix such as fibronectin or collagens, and in HIV infection.
Species Reactivity:	Reacts with human. Others not known.
Positive Control:	U-87 MG (Human glioblastoma astrocytoma), Ramos (Human Burkitt's lymphoma, HEP-G2 cells, and lymphocytes. Lymph nodes and tonsils.
Cellular Localization:	Cell Surface.
Titer/ Working Dilution:	Immunohistochemistry (Frozen): 0.5-1 µg/ml Flow Cytometry: 0.5-1 µg/million cells Immunofluorescence: 0.5-1 µg/ml
Microbiological State:	This product is not sterile.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

**CE**

EC | REP  
Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands

**Uses/Limitations:** Not to be taken internally.  
For Research Use Only.  
This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.  
Do not use if reagent becomes cloudy.  
Do not use past expiration date.  
Non-Sterile.

**Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)**

**Procedure:**

1. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
2. **Visualization:** For maximum staining intensity we recommend the “CRF Anti-Polyvalent HRP Polymer (DAB) Lab Pack” (ScyTek catalog# CPP125, see IFU for instructions), combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).


**Precautions:** Contains Sodium Azide as a preservative (0.09% w/v).  
Do not pipette by mouth.  
Avoid contact of reagents and specimens with skin and mucous membranes.  
Avoid microbial contamination of reagents or increased nonspecific staining may occur.  
This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


**References:**

1. Fleischer B. CD26: a surface protease involved in T-cell activation. Immunol Today 1994, 15(4):180-184.
2. Abbott CA. et. al. Genomic organization, exact localization, and tissue expression of the human CD26 (dipeptidyl peptidase IV) gene. Immunogenetics 1994, 40:331-338.
3. Valenzuela A et. al. HIV-1 envelope gp120 and viral particles block adenosine deaminase binding to human CD26. Adv Exp Med Biol 1997, 421:185-192.
4. Cheng HC, et al. Lung endothelial dipeptidyl peptidase IV promotes adhesion and metastasis of rat breast cancer cells via tumor cell surface-associated fibronectin. J Biol Chem. 1998 273(37):24207-24015.

**Warranty:**

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.



EC REP

Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands